

Approximate zeta functions for the Sinai billiard and related systems

Per Dahlqvist

Mechanics Department

Royal Institute of Technology, S-100 44 Stockholm, Sweden

Abstract - We discuss zeta functions, and traces of the associated weighted evolution operators for intermittent Hamiltonian systems in general and for the Sinai billiard in particular. The intermittency of this billiard is utilized so that the zeta functions may be approximately expressed in terms of the probability distribution of laminar lengths. In particular we study a one-parameter family of weights. Depending on the parameter the trace can be dominated by branch cuts in the zeta function or by isolated zeros. In the former case the time dependence of the trace is dominated by a powerlaw and in the latter case by an exponential. A phase transition occurs when the leading zero collides with a branch cut. The family considered is relevant for the calculation of resonance spectra, semiclassical spectra and topological entropy.

1 Introduction

The space of Hamiltonian systems is framed by two extreme cases - integrable systems and strongly chaotic systems, the latter in the sense of Axiom-A. Both cases may be successfully *solved*, but the methods of solution look entirely different. For integrable systems *solving* means *finding* the constants of motion. In the Axiom-A case one rather aims at a statistic description of the motion - one calculates different kinds of entropies, fractal dimensions, resonance spectra etc. To this end *cycle expansions* of zeta functions have shown very successful [1, 2, 3]. So far, the success of this approach has been limited to the Axiom-A case. The reason is that Axiom-A ensures nice analytic features of the zeta functions involved [4, 5, 6, 7], whereas different kinds of singularities emerges when departing from Axiom-A [8]. If the system is not Axiom-A, but still ergodic, it is generally intermittent (there are cycles with arbitrary small Lyapunov exponents) and lacks a simple symbolic dynamics. This is e.g. the case for most bound ergodic systems [9]. It is of great interest to extend the study of zeta functions to this case. The resulting theory would be an analogue of perturbation theory for almost integrable systems.

Such a theory would be interesting not only for ergodic systems. The motion in a chaotic sea of a generic mixed system is in general intermittent. This is because the trajectory may be trapped inside *cantori* surrounding the stable islands, where it exhibits quasi-integrable motion.

The cycle expansion for a system with a finite subshift symbolic dynamics consists of two parts [1], a *fundamental part*, giving the gross structure of the spectrum, and *curvature corrections*.

When the symbolic dynamics is an infinite subshift, there is no similar division. One goal of this paper is to establish a fundamental part for this case. One can think of two ways out

of the dilemma. One can define successive finite subshifts, approaching the infinite subshift, by utilizing a *pruning front* [10, 11]. This mean that one persists on a periodic orbit description. We will attack the problem from the opposite end and completely abandon the periodic orbits. We will thus make extensive use of our systems being ergodic and Hamiltonian, so that the invariant density is known a priori. One of the morals to be taught is that there is no reason to find a lot of periodic orbits in order to compute something you already know in advance. The idea to use probabilities rather than periodic orbits has been used in a similar way in ref. [12].

In section 2 we develop most of the theoretical apparatus needed. The exposition will be rather brief as most of the material can be found elsewhere. In section 3, we apply these ideas to the Sinai billiard and formulate an approximate zeta function. In section 4 we study this zeta function numerically and compare with periodic orbit theory.

2 Theory

Much of the theoretical work on these matters are centred around the evolution operator. It describes the evolution of a phase space density $\Phi(x)$

$$\mathcal{L}_w^t \Phi(x) = \int w(x, t) \delta(x - f^t(y)) \Phi(y) dy \quad . \quad (1)$$

The phase space point x is taken by the flow to $f^t(x)$ during time t . $w(x, t)$ is a weight associated with a trajectory starting at x and evolved during time t . It is multiplicative along the flow, that is $w(x, t_1 + t_2) = w(x, t_1) w(f^{t_1}(x), t_2)$.

We emphasize that t is a continuous variable as we are studying flows and not maps.

2.1 Trace formulas and zeta functions

Our main concern in this paper is to compute the trace of the evolution operator, that is, the sum of its eigenvalues. The material in this subsection may be found e.g. in refs. [1, 13]

The trace may be written as a sum over the periodic orbits in the system

$$\begin{aligned} \text{tr} \mathcal{L}_w^t &= \int w(x, t) \delta(x - f^t(x)) dx = \\ &= \sum_p T_p \sum_{n=1}^{\infty} w_p^n \frac{\delta(t - nT_p)}{|\det(1 - M_p^n)|} \quad , \end{aligned} \quad (2)$$

where n is the number of repetitions of primitive orbit p , having period T_p , and M_p is the Jacobian of the Poincaré map. w_p is the weight associated with cycle p .

The trace is written as the Fourier transform of the logarithmic derivative of a *zeta function*

$$\text{tr} \mathcal{L}_w^t = \frac{1}{2\pi i} \int_{-\infty}^{\infty} e^{ikt} \frac{Z'_w(k)}{Z_w(k)} dk \quad . \quad (3)$$

We restrict ourselves to systems with two degree's of freedom for which all periodic orbits are isolated and unstable. The zeta function then reads

$$Z_w(k) = \prod_p \prod_{m=0}^{\infty} \left(1 - w_p \frac{e^{-ikT_p}}{|\Lambda_p| \Lambda_p^m} \right)^{m+1} \quad , \quad (4)$$

where Λ_p is the expanding eigenvalue of M_p .

Putting $w = 1$ we obtain a zeta function whose zeros yields the so called *resonance spectrum* or spectrum of correlation exponents,

$$Z(k) = \prod_p \prod_{m=0}^{\infty} \left(1 - \frac{e^{-ikT_p}}{|\Lambda_p| \Lambda_p^m} \right)^{m+1} \quad . \quad (5)$$

The leading zero, k_0 , is the escape rate, which is zero for a bound system.

By using different weights w one can probe different properties of the flow.

The topological entropy is obtained by considering the weight $\omega = |\Lambda(x, t)|$. $\Lambda(x, t)$ is the expanding eigenvalue of the Jacobian transverse to the flow. It is only approximately multiplicative along the flow but it is possible to modify it slightly so as to become exactly multiplicative [14]. However, this subtle difference is of no importance for our purposes. The trace formula then becomes

$$\text{tr} \mathcal{L}_{top}^t \approx \sum_p T_p \sum_{n=1}^{\infty} \delta(t - nT_p) . \quad (6)$$

The leading zero lies on the negative imaginary axis. The topological entropy is then $h = ik_0$, the asymptotic behaviour of the trace is

$$\sum_p T_p \sum_{n=1}^{\infty} \delta(t - nT_p) \rightarrow e^{ht} \quad (7)$$

so that the number of cycles with periods less than t is $\sim e^{ht}/ht$, which is the familiar result.

The semiclassical case is obtained by considering the weight $w = \sqrt{|\Lambda(x, t)|}$. We now get the following zeta function,

$$Z_{sc}(k) = \prod_p \prod_{m=0}^{\infty} \left(1 - \frac{e^{-ikT_p}}{\sqrt{|\Lambda_p|} \Lambda_p^m} \right)^{m+1} . \quad (8)$$

This zeta function is called *the quantum Fredholm determinant* [3] and is equivalent to the Gutzwiller-Voros [15, 16] zeta function in the semiclassical limit, and have indeed nicer analytical features. We have left out the Maslov indices but it is possible to account for them in the weight as well.

2.2 Symmetry decomposition of $Z(k)$

Many systems possess finite symmetries. This enables a factorization of the zeta function. This was first shown for the semiclassical case in refs. [17, 18] and for a more general setting in ref. [19]. The zeta function $Z_w(k)$ in (4) is then written as a product $Z_w(k) = \prod_r Z_{w,r}(k)$ over r , the irreducible representations of the symmetry group G , and each $Z_{w,r}$ is given by

$$Z_{w,r}(k) = \prod_p \prod_{m=0}^{\infty} \left(1 - \chi_r(g_p) w_p \frac{e^{-iT_p k}}{|\Lambda_p| \Lambda_p^m} \right)^{m+1} , \quad (9)$$

where p runs over all prime cycles in the fundamental domain. $\chi_r(g_p)$ is the group character for the irreducible representation r , and g_p is the group element obtained as the product of group factors associated with the reflections of the domain walls.

Throughout the paper we will restrict ourselves to one-dimensional irreducible representations of the group. Periodic orbits running along symmetry lines must be given a special treatment. We will discuss this when the result is needed.

In the quantum case the different irreducible representations corresponds to the different symmetry classes of the wave functions. But in the general classical case it is harder to give a comprehensible interpretation. The important thing to keep in mind in the following is that the leading zero is always in the symmetric representation.

2.3 Zeta functions in the BER approximation

In ref. [8] an approximate expression for the zeta function is given for intermittent, ergodic Hamiltonian systems. The idea is based on a paper by Baladi, Eckmann and Ruelle [20] and it is therefore referred to as the BER approximation.

In an intermittent system there are two, more or less, distinct phases; one regular and one irregular (chaotic). Call the consecutive instants when the system enters the regular phase $\{t_i\}$ and consider the intervals $I_i = [t_{i-1}, t_i]$. Provided the chaotic phase is *chaotic enough*, the motions in different intervals are nearly mutually independent. This is called *assumption A*. In particular, the lengths of these intervals $\Delta_i = t_i - t_{i-1}$ are mutually independent and Δ may be considered as a stochastic variable with probability distribution $p(\Delta)$. Under this assumption, the zeta functions may be expressed in terms of the Fourier transform of $p(\Delta)$

$$Z(k) \approx \hat{Z}(k) \equiv 1 - \int_0^\infty e^{-ik\Delta} p(\Delta) d\Delta \quad (10)$$

We use the $\hat{\cdot}$ -symbol to denote all quantities in the BER approximation.

This approximation is not restricted to unit weight. If the weight associated with one interval is a function of the length of the interval Δ only, then

$$\hat{Z}_w(k) = 1 - \int_0^\infty e^{-ik\Delta} w(\Delta) p(\Delta) d\Delta \quad . \quad (11)$$

This is one of the central ideas of this paper. To show this, one just repeats the calculation of ref. [8], squeezing in the weight at the appropriate places. As it is straightforward to do we will not perform it here. The assumption above, that the weight is a function of Δ only, is quite reasonable for many systems. For the Sinai billiard, which we will study soon, the expression has to be slightly generalized.

3 The Sinai billiard

The purpose of this section is to apply the BER approximation, that is formulate $\hat{Z}_{w,r}(k)$, for the Sinai billiard [21] and compare with cycle expansions. The billiard consist of a unit square with a scattering disk, having radius R ; $0 < 2R \leq 1$, centered on its midpoint, cf. fig. 1b. The Sinai billiard is fairly simple but have all the typical features for bound chaotic system; it is intermittent and lacks a simple symbolic dynamics. Moreover, it has marginally stable orbits. However, a trajectory in the hyperbolic phase (hyperbolic in the sense that all cycles there are unstable and isolated) will never end up on a marginally stable orbit, it can only come arbitrary close to. This means that we can separate out this marginal phase. That is, the integral (2) is only performed over the hyperbolic part of phase space, thus excluding a region of measure zero, so that the sum in eq. (2) runs only over isolated periodic orbit.

This separation cannot be performed in the quantum case due to an extra contribution from the marginal orbits only disappearing in the limit $E \rightarrow \infty$. The study of the quantum case will be postponed to a forthcoming paper.

The trajectory of the Sinai billiard consists of laminar intervals, when bouncing between the straight sections, interrupted by scatterings on the central disk. The Sinai billiard seems therefore ideally suited for the BER approximation. When the disk radius is small, the memory of the previous laminar interval should be almost completely lost. We will not discuss corrections to the BER approximation arising from the correlations between laminar intervals. We simply assume that they are small when the disk radius is small. The length of the chaotic interval is infinitely short so that Δ is simply the length of the trajectory between two disk scatterings. For big radii, there is another source of intermittency [22, 23], the trajectory may be trapped between the disk and the straight section. For these reasons we will focus on the limit of small disk radii.

3.1 Symbolic dynamics

First we will define a symbolic dynamics for the Sinai billiard. The reason is twofold. First we need it for the application of the BER approximation. Secondly we will use it for finding the periodic orbits of the system, which we will need to test the results obtained from the BER approximation.

Figure 1: Three equivalent representations of the Sinai billiard. a) The fundamental domain. b) The original Sinai billiard with definitions of the variables ϕ and α . c) The unfolded system with a sample of orbits which are periodic of period one in the fundamental domain.

We let the disk be our Poincaré surface of section. The canonical variables are $(\xi, p_\xi) = (R\phi, \sqrt{2E}\sin\alpha)$ where the angles ϕ : $0 < \phi < 2\pi$ and α : $-\frac{\pi}{2} < \alpha < \frac{\pi}{2}$ are defined in fig. 1b. The area preserving map $(\xi, p_\xi) \mapsto (\xi, p_\xi)$ has uniform invariant density.

We now want to define a symbolic dynamics, such that each iterate of the map corresponds to one symbol, that is, each symbol corresponds to one laminar interval.

To do this we define two equivalent billiard systems, derived from the original system. The original billiard has the symmetry group $G = C_{4v}$, [24]. This group has eight elements in five conjugacy classes, see table 1.

The first derived system is the fundamental domain, or the desymmetrized system, see fig 1a. This system has, by definition, no symmetry, except for time reversal.

To obtain the second derived system we go in the opposite direction. We unfold the billiard, see fig 1c, into a regular lattice of disks. Apart from the group C_{4v} this system has also a discrete translational symmetry.

We note that any trajectory in the unfolded billiard may be encoded by a sequence of column vectors $\dots Q_{i-1} Q_i Q_{i+1} \dots$

$$Q_i \in \Omega$$

$$\Omega : \left\{ \begin{pmatrix} n_x \\ n_y \end{pmatrix}; n_x, n_y \in \mathbb{Z}, \gcd(n_x, n_y) = 1 \right\} \quad (12)$$

where $\gcd(n_x, n_y)$ stands for the greatest common divisor of n_x and n_y . The number n_x (n_y) simply tells the number of disk sites the segment of the trajectory has travelled in the horizontal (vertical) direction. It is easy to see that non coprime Q 's cannot be realised.

A general trajectory corresponds to a semi infinite string of Q 's. We will focus on periodic strings $\overline{Q_1 Q_2 \dots Q_n}$. Such a string uniquely determines the trajectory in the unfolded billiard up to a translation which means that the corresponding trajectory is completely determined in the original billiard, and is indeed periodic there. Of course, some trajectories defined in this way would need to go through disks. The corresponding symbol sequences are then said to be *pruned*. But when the disk radius is small we expect this pruning to apply only to those orbits with some *long* segment(s). More exactly if all the Q 's have $\sqrt{n_x^2 + n_y^2} \ll 1/R$ we expect the pruning to be negligible. This is very convenient for computing purposes since a large fraction of the generated symbol sequences will be realised as true orbits.

We observe that a symmetry transformation of the orbit corresponds to a transformation $Q_i \mapsto gQ_i$ where g is an element of $C_{4,v}$ represented by a 2×2 matrix, see table 1.

Our goal is to encode cycles in the fundamental domain. Due to symmetry, several cycles of Q 's correspond to one and the same cycle in the fundamental domain. We therefore want to translate the cyclic symbol string $\overline{Q_1 Q_2 \dots Q_n}$ to another string $\overline{s_1 s_2 \dots s_n}$ where the s 's are taken from another alphabet. The translation must fulfill the following conditions:

1. The translation is unique.
2. The sequence $\overline{s_1 s_2 \dots s_n}$ is invariant under a symmetry operation on $\overline{Q_1 Q_2 \dots Q_n}$.
3. A cyclic shift on the Q 's corresponds to a cyclic shift on the s 's.

This will be (almost) achieved in the following way. Let each s_i be an ordered pair $s_i = (q_i, g_i)$

$$\begin{aligned} g_i &\in C_{4,v} \\ q_i &\in \omega \end{aligned} \quad (13)$$

$$\omega : \left\{ \begin{pmatrix} n_x \\ n_y \end{pmatrix}; n_x, n_y \in \mathbb{N}, n_x \geq n_y, \gcd(n_x, n_y) = 1 \right\}$$

The translation is defined by:

$$\begin{aligned} Q_1 &= g_0 g_1 q_1 \\ Q_2 &= g_0 g_1 g_2 q_2 \end{aligned} \quad (14)$$

E	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	Identity
I	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	Inversion
R^+, R^-	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	Rotation in the positive and negative directions
σ_+, σ_-	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Reflection in the vertical and horizontal axis
$\sigma_/, \sigma_\backslash$	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Reflections in the diagonals

Table 1: The elements of the group $C_{4,v}$. They are represented by 2×2 matrices operating on the column vectors q_i and Q_i .

$$\begin{aligned}
& \vdots \\
Q_n &= g_0 g_1 \dots g_n q_n
\end{aligned} \tag{15}$$

The translation is unique as long as no vector q_i is equal to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ or $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ (we refer to these as boundary symbols). We will deal with this problem in a while. Condition 2 is also fulfilled - a symmetry operation affects only g_0 . It is easy to check that also condition 3 is fulfilled (a cyclic shift also affects g_0).

Suppose now that q_j is a boundary vector. If we infer the restriction that g_j must not be a reflection ($\sigma_+, \sigma_-, \sigma_/$ or $\sigma_\backslash$) the translation is again unique provided that not *all* the q 's are boundary symbols.

Suppose the latter is the case and that we have found *one* translation of the string $\overline{Q_1 Q_2 \dots Q_n}$, namely: $\overline{s_1 s_2 \dots s_n}$. It is now possible to find another s -string corresponding to the same Q -string. This is done in the following way. Suppose that q_i has the symmetry ϕ_i , that is, if $q_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ then $\phi_i = \sigma_-$ and if $q_i = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ then $\phi_i = \sigma_/$. Then the transformed sequence

$$\begin{aligned}
q_i &\mapsto q_i \\
g_i &\mapsto \phi_{i-1} g_i \phi_i \quad i \neq 0 \\
g_0 &\mapsto g_0 \phi_n
\end{aligned} \tag{16}$$

corresponds to the same Q string. Remember that the string is periodic so that e.g. $\phi_0 = \phi_n$. Performing this transformation once more gives back the original s string. So there remains a problem that an orbit in the fundamental domain might correspond to two symbol strings and we have to account for this problem when generating the periodic orbits. The symbolic dynamics defined is still a good one and the problem above should be interpreted as a boundary effect.

As we said, the problem arises only if all vectors are on the boundary. If one is not, and we try to perform the transformation (17), we see that at least one g_i corresponding to a boundary q_i would transform to a reflection and such strings have already been forbidden.

The group characters in eq. (9) are determined both from the disk bounces and the bounces on the square walls. They are easily expressed in terms of the symbol code

$$\chi_r(g_p) = \prod_{i=1}^{n_p} \chi_r(g_i) \chi_r(\sigma_-)^{n_{x,i} + n_{y,i}} \quad , \tag{17}$$

To obtain this we have used the familiar property of group characters: $\chi_r(g_1 g_2) = \chi_r(g_1) \chi_r(g_2)$. where g_i , $n_{x,i}$ and $n_{y,i}$ are taken from the symbols string corresponding to prime orbit p .

One may also consider the Sinai billiard with periodic boundary conditions. One then simply exclude the reflection factor $\chi_r(\sigma_-)$ in eq. (17).

3.2 Free Directions

For any disk radius $0 < 2R < 1$ there are a finite number of directions through which a trajectory may go without ever bouncing on any disk. We call them *free directions*. Consider the direction vector (x, y) . A direction cannot be free if x/y is irrational. So we may take (x, y) as coprime integers such that $x/y \leq 1$. The direction (x, y) is found to be free if

$$2R < \frac{1}{\sqrt{x^2 + y^2}} . \quad (18)$$

To show this, consider an integer lattice of points (or consult ref [28]). We are seeking the lattice point (ξ, η) closest to the line from $(0, 0)$ to (x, y) and lying below. The closest point above this line is then, due to symmetry, $(x - \xi, y - \eta)$. By the definition of Farey sequences, $\frac{\eta}{\xi}$ is the preceeder of $\frac{y}{x}$ in the Farey sequence of order x . The Farey theorem then says that $\xi y - \eta x = 1$ [25]. This is nothing but the area of the parallelogram spanned by the points $(0, 0)$, (ξ, η) , (x, y) and $(x - \xi, y - \eta)$ so that the distance between the point (ξ, η) to the diagonal is $1/\sqrt{x^2 + y^2}$ and the result above follows.

The direction (x, y) being free is equivalent with the existence of the periodic orbit $\bar{s}$ where $s = ((\frac{x}{y}), E)$.

3.3 The zeta function in the BER approximation

We begin by considering the symmetric representation $r = A_1$ (then $\chi_r(g_p) = 1$) and unit weight $w = 1$ in the zeta function. The corresponding indices will be omitted in the zeta function.

Utilizing the uniformity of the invariant measure [21] we can write down the following expression for the probability distribution $p(\Delta)$

$$p(\Delta) \approx \sum_{q \in \omega} a_q(0) \delta(\Delta - T_q) , \quad (19)$$

where $a_q(0)$ as given by

$$a_q(0) = \frac{2}{\pi} \int_{\Omega_q} d\phi d(\sin \alpha) . \quad (20)$$

Ω_q is the part of phase space (i.e. the α, ϕ -plane) for which the trajectory hit disk q (in the unfolded system). We have used the symmetry by letting $q \in \omega$, meaning that we take only orbits going out into the first octant into account. Only the fractions of the disks $(\frac{1}{0})$ and $(\frac{1}{1})$ lying in this octant are included. The argument (now put to zero) in $a_q(0)$ is introduced for later purposes.

We have made the approximation that all trajectories to disk q have the same length l_q corresponding to the time of flight T_q (this will be refined a little in section 3.5). This is reasonable when the disk radius is small. Taking the Fourier transform we get:

$$\hat{Z}(k) = 1 - \sum_q a_q(0) e^{-ikT_q} . \quad (21)$$

Although we use equality sign above we must keep in mind that this result involves an approximation in addition to the BER approximation.

3.4 Relation to cycle expansion of $Z(k)$.

We are now interested in the relations between the approximate zeta function $\hat{Z}(k)$ and the cycle expansion of the exact zeta function $Z(k)$. First we elaborate a little more on $\hat{Z}(k)$.




Figure 2: Comparison of the prefactors of eqs (22) and (25) plotted versus $\sin\phi$

It is convenient to write $a_q(0) = \hat{\eta}_q \frac{8R}{\pi l_q}$, where $\frac{8R}{\pi l_q}$ is the normalized phase space area taken up by disk q if it is not shadowed, and $\hat{\eta}_q$ is a pruning factor; $0 \leq \hat{\eta}_q \leq 1$. We thus write

$$\hat{Z}(k) = 1 - \sum_q \hat{\eta}_q \frac{8R}{\pi l_q} e^{-ikT_q} . \quad (22)$$

The direction $(1,0)$ is always free if $2R < 1$. This means that every disk $\binom{n}{1}$ is accessible and takes up a normalized phase space area

$$a_{q=\binom{n}{1}}(0) \approx \frac{2}{\pi} (2R + 1/2R - 2) \frac{2}{n(n-1)(n-2)} \quad (23)$$

Let's now consider the zeta function $Z(k)$. First we will neglect the factors with $m > 0$ in (5). Given a periodic orbit $\overline{s_1 s_2 \dots s_n}$ we make the following assumption (*assumption B*):

1. The accessible values for s_i does not depend on s_j , $j \neq i$.
2. The stability eigenvalues are given by $\Lambda_{\overline{s_1 \dots s_n}} = \prod_i \Lambda_{\overline{s_i}}$
3. The periods are given by $T_{\overline{s_1 \dots s_n}} = \sum_i T_{\overline{s_i}}$

This assumption is not at all true, although it is realistic if all $T_{\overline{s}}$'s are small. We introduce it as a working assumption to shed some light on the nature of the BER approximation.

Condition 1 means that we can make a curvature expansion [1]. Conditions 2 and 3 means that all curvatures disappear and what remains is the fundamental part

$$Z(k) = 1 - \sum_s \frac{\eta_{\overline{s}}}{\Lambda_{\overline{s}}} e^{-ikT_{\overline{s}}} . \quad (24)$$

$\eta_{\overline{s}}$ is a pruning factor which is $= 1$ if cycle $\overline{s}$ exist, and zero otherwise. Since we have assumed that $T_{\overline{s=(q,g)}}$ is a function of q only ($\equiv T_q$) we sum over $g \in C_{4,v}$ keeping $q = \binom{n_x}{n_y}$ fixed and get

$$Z(k) = 1 - \sum_{q \in \omega} \eta_q \frac{R}{2l_q} \left[(1 + \sqrt{2})(1 + \cos\phi_q) + \sin\phi_q \right] e^{-ikT_q} . \quad (25)$$

The angle ϕ_q is the angle between the horizontal axis and the direction vector (n_x, n_y) . η_q takes on discrete values and is $= 1$ if all g 's are realised. In fig. 2 we compare the prefactors (in arbitrary units) of eqs. (22) and (25) provided that $\eta_q = \hat{\eta}_q$. The similarity with eq. (24) is obvious and we may deduce that Assumptions A and B are similar. The reason for the agreement is that the available phase space region in eq. (21) is extrapolated from the local stabilities in eq. (25). This should work well as long as there is no shadowing (=obstructing) disks in between.

We will now see that eq. (25) fails completely when pruning becomes essential, that is when $\hat{\eta}_q \ll 1$. This is e. g. the case far out in the free directions. Consider the free direction $(1, 0)$. We see from (23) that $a_{q=(\begin{smallmatrix} n \\ 1 \end{smallmatrix})}(0)$ decays as $\sim 1/n^3$. There is only one period-one cycle with $q = (\begin{smallmatrix} n \\ 1 \end{smallmatrix})$, and it is the one having $g = \sigma_-$ and the corresponding prefactors $1/\Lambda_{(q, \sigma_-)}$ decays as $1/n^2!$

The reason for the discrepancy is the fake assumption B. An apparent example of violation of B1 is the following. The cycle $\bar{s}$ where $s = (q, g = E)$ is pruned but the same symbol s may very well appear in longer cycles.

As we have already said, there is no well defined fundamental part of the expanded zeta function. Indeed, by our discussion we are led to suggesting that the BER approximation provides a natural generalization of the fundamental part, with the extra advantage of preserving unitarity: $k_0 = 0$ is by construction a zero of the unweighted zeta function $\hat{Z}(k)$.

We have seen that the problem of establishing a fundamental part of the zeta function using periodic orbits is connected with the marginal orbits which we have pruned by excluding the marginal phase and the accumulation of hyperbolic orbits towards it. The BER approximation tells us that there may be simpler means of exploring the available phase space for a certain symbol than letting periodic orbits explore it.

3.5 Other representations, other weights in the BER approximation

It is straightforward to study other than the symmetric representation A_1 in the BER approximation. In this approximation we only consider one single disk bounce. Since we have decided that the trajectory goes out into the first octant it suffices to determine what octant it came from in order to determine the group element g_s associated with this particular bounce. That is, we divide the phase space region Ω_q into eight regions Ω_s where $s = (q, g)$ and $g \in C_{4,v}$. Some of these may of course be empty.

The group character associated with the whole interval (one bounce + the trajectory to the next disk) is then (cf, (17))

$$\chi_r(g_s) = \chi_r(g)\chi_r(\sigma_-)^{n_x+n_y} . \quad (26)$$

The zeta function may then be written

$$\hat{Z}_r(k) \approx 1 - \sum_{s=(q,g)} \chi_r(g_s) a_s(0) e^{-ikT_s} , \quad (27)$$

where in $a_s(0)$ is defined by (20) but the integral extends only over Ω_s .

It is also straightforward to use other weights w in the BER approximation for the Sinai billiard. All weights we are going to study in this paper belong to the family $w = |\Lambda(\phi, \alpha)|^\tau$. The corresponding family of zeta functions then reads

$$\hat{Z}_{\tau,r}(k) \approx 1 - \sum_s \chi_r(g_s) a_s(\tau) e^{-ikT_s} , \quad (28)$$

where

$$a_s(\tau) = \frac{2}{\pi} \int_{\Omega_s} |\Lambda(\phi, \alpha)|^\tau d\phi d(\sin \alpha) \quad (29)$$

$$\Lambda(\phi, \alpha) \approx \frac{2l_s}{R \cos \alpha} . \quad (30)$$

As $\Lambda(\phi, \alpha)$ is not constant over Ω_s , c.f. eq. (11), we have averaged it over Ω_s .




Figure 3: The trace of the unweighted evolution operator. The parameters are $R = 0.1$, $r = A_1$, $\sigma = 0.1$

4 Computations and results

We now turn to explicit computations. The aim is to check the validity of the BER approximation as outlined in section 3.5, meaning that we restrict ourselves to the family of weights $w = |\Lambda|^\tau$.

The strategy is the following. First, we compute the trace in the BER approximation,

$$\text{tr} \hat{\mathcal{L}}_{\tau,r,\sigma}^t = \frac{1}{2\pi i} \int_{-\infty}^{\infty} e^{ikt} \frac{\hat{Z}'_{\tau,r}(k)}{\hat{Z}_{\tau,r}(k)} e^{-k^2 \sigma^2 / 2} dk . \quad (31)$$

Then, in order to check this result, we compute the 'exact' trace, by means of the trace formula

$$\text{tr} \mathcal{L}_{\tau,r,\sigma}^t = \sum_p T_p \sum_{n=1}^{\infty} \chi_r(g_p)^n |\Lambda_p|^{n\tau} \frac{e^{(t-nT_p)^2 / 2\sigma^2}}{|(1 - \Lambda_p^n)(1 - 1/\Lambda_p^n)|} , \quad (32)$$

To this end we need the periodic orbits. We use gaussian smearing in order to avoid the delta peaks.

The full trace of the full system (fig 1b) is obtained as a sum of traces for all irreducible representations of $G = C_{4,v}$. As we will not treat two dimensional representations we will not perform the sum.

We mentioned before that orbits running along symmetry lines has to be given a special treatment. There are two such cycles in the Sinai billiard. We treat them in the following way. One has symmetry σ_- . We simply exclude from the trace formula (32) if $\chi_r(\sigma_-) = -1$. The other one, having symmetry σ_+ is treated analogously. This procedure is not exact, c. f. [18, 19], but very accurate.

The computational task lies in determining the quantities $a_s(\tau)$ as defined by eq. (29).

We do this in the simplest thinkable way by binning the phase space corresponding to a disk q and deleting the phase space shadowed by nearer disks. More exactly, we bin the α variable, keeping track of the limits in the ϕ direction.

The integrals will for obvious reasons diverge if $\tau > 2$.

Of course, we only calculate a finite number of a_s . Then there are infinite tails in all free directions. Let us for a minute restrict ourselves to the case $r = A_1$ and the $(1,0)$ free direction. Far out in the free directions only a small fraction of disk $q = \begin{pmatrix} n \\ 1 \end{pmatrix}$ is visible. Then $l_{s=(q,g)}$ is essentially independent of g . If we sum over g keeping q fixt

$$\sum_g a_{s=\begin{pmatrix} n \\ 1 \end{pmatrix},g}(\tau) \approx \frac{2}{\pi} \left(\frac{2n}{R}\right)^\tau \frac{(2/R-4)^{1-\tau/2}(1-2R)}{2-\tau} \frac{1}{n^{1+\tau/2}(n-1)^{1-\tau/2}(n-2)^{1-\tau/2}} \quad (33)$$

We will not bore the reader with details about the other r 's and free directions. These infinite tails will make the zeta functions diverge in the upper half k -plane. It is thus crucial that we find an analytical continuation of the zeta function.

We also have to choose the lengths l_s associated with Ω_s . We simply choose it to be the period of the cycle $\bar{s}$ regardless if it is pruned or not. We choose the energy such that the lengths l_s and the times T_s coincides.

We will also calculate all periodic orbits up to some maximal period. We thus generate symbol codes according to the scheme outlined in the previous section. The orbits are then found by an extremum principle according to ref. [26]. Last it is checked if each cycle is allowed or pruned.

4.1 The unweighted zeta function, $\tau = 0$.

We begin by considering the summation of the tails. Consider disk $q = \begin{pmatrix} n \\ 1 \end{pmatrix}$ in the free direction $(1,0)$. Only a small fraction of this disk is visible for large n . Then $\Omega_{s=(q,g)}$ is nonempty only for $g = E$ and $g = \sigma_-$ and all points lie on about the same distance $l_s = l_q$. Restricting ourselves to $r = A_1$ we have to consider the following series

$$\sum_{n=N+1}^{\infty} \frac{2}{n(n-1)(n-2)} e^{-ikT_q} . \quad (34)$$

If we as an approximation put $l_{q=(n,1)} \approx n$ the series reduces to a Fourier series in $z = \exp(-ik)$. This is easily summed since

$$\sum_{n=3}^{\infty} \frac{z^n}{n(n-1)(n-2)} = -\frac{1}{2}(1-z)^2 \log(1-z) + \frac{3}{4}(1-z)^2 - (1-z) + \frac{1}{4} . \quad (35)$$

The zeta function will thus have branch cuts along $Re(k) = 2\pi N$ and $Im(k) > 0$.

There are similar contributions in all free direction. The free direction (x,y) will induce branch cuts along $Re(k) = 2\pi N/\sqrt{x^2+y^2}$ and $Im(k) > 0$.

Close to the origin $k = 0$ the zeta function behaves as $Z(k) = c_1(ik) + d_2(ik)^2 \log(ik) + c_2(ik)^2 \dots$, ($r = A_1$). Consequently the trace will have the asymptotic behaviour


$$tr \hat{\mathcal{L}}_{\tau=0, r=A_1, \sigma}^t \sim 1 - C/t \quad (36)$$

where C is a positive constant, it may be related to the expectation value of $p(\Delta)$ [8]. The convergence of the trace formula towards unity is usually referred to as a periodic orbit sum rule [27].


When considering other representations, r , some tail series will have alternating signs. This will shift the branch cuts in the real direction. These zeta functions will not have a root at the origin and the traces go asymptotically to zero.

We now turn to the numerical computation of the trace. When we calculate the Fourier transform we simply make the integration along the the real axis, except at the origin which we sneak just below.


With the number of periodic orbits calculated (between 5000 and 8000 depending on R) we cannot check the asymptotic limit discussed above. To do the numerical check we restrict ourselves to much shorter times t . The result is seen in figs 3 and 4. The agreement is in most cases very good. We note




a) $R = 0.2, r = B_2, \sigma = 0.1$



b) $R = 0.2, r = A_2, \sigma = 0.1$



c) $R = 0.1, r = B_2, \sigma = 0.1$



d) $R = 0.1, r = A_2, \sigma = 0.1$

Figure 4: Trace of the unweighted evolution operator obtained from the BER approximation and directly from periodic orbits

- For small t the agreement is less good or even lousy. The exact result for small t depends on a small number of cycles and the BER theory requires an average effect from many cycles.
- For fixed representation r , the agreement is better for small radii R . This is exactly what we expected.
- For fixed radius R the agreement is best for the symmetric representation A_1 and worst for the anti symmetric A_2 . In the former case all Ω_s provides the same sign but otherwise different phase space regions come with different signs (provided by the group characters χ). It takes more periodic orbits in order to provide averages for this finer partition.

We have seen that the trace is almost entirely governed by the branch cuts. (There are a few isolated zeros but they are extremely sparse and have no practical importance). We are thus faced with an entirely different situation than for Axiom-A systems for which the zeta function is entire and the zeros gives the discrete spectrum of the operator.

We have not discussed in what sense the cuts may be interpreted as a continuous part of the spectra. If the trace, by means of the sum over eigenvalues, is not mathematically well defined, we let it be defined by means of the corresponding periodic orbit sum.

4.2 Topological entropy, $\tau = 1$

We now have to find the analytic continuation for sums of the form, cf. eq. (33),

$$\sum_{n=N+1}^{\infty} \frac{e^{-ikcn}}{n^{3/2}} . \quad (37)$$

This is just a power series in $\exp(-ikc)$. We observe that

$$\sqrt{1-z} = 1 - \sum_{n=1}^{\infty} \frac{(2n-3)!!}{(2n)!!} z^n . \quad (38)$$

Using Stirlings formula we find the following asymptotic expression for the coefficient

$$\frac{(2n-3)!!}{(2n)!!} \sim \frac{1}{\sqrt{4\pi}} \frac{1}{n^{3/2}} \quad (39)$$

enabling an summation of the (37), provided of course that N is sufficiently big. The zeta function still have cuts in the upper half plane. However, there are isolated zeros far down in the lower half plane. The determination of these does not depend very much on the summation of the tails. Therefore we could take only the leading order of the tails series into account above.

When performing the integral (31) we let the contour go below all singularities of the integrand, that is below all discrete zeros k_α of $\hat{Z}$. The sum over residues gives


$$tr \hat{\mathcal{L}}_{\tau=1, r=A_1, \sigma}^t \sim \sum_{\alpha} e^{ik_\alpha t} e^{-k_\alpha^2 \sigma^2 / 2} . \quad (40)$$

The result is seen in fig. 5. This result is again good for large t .

Of course, in the contour integration we have neglected the contribution coming from the integration around the branch cuts cf ref. [8]. This should give an extra contribution $\sim -1/\sqrt{t}$ which is of course overwhelmed by the exponential increase, but could explain the discrepancy for small t in fig. 5.

We have determined the topological entropy $h(R)$ for two different radii R . We find $h(0.2) = 2.1346$ and $h(0.1) = 2.02024$. The entropy $h(R)$ tends to a finite limit when $R \rightarrow 0$. Its limiting value is obtained from the equation

$$1 - \sum_{i,j \in \mathbb{Z}, \gcd(i,j)=1} e^{-h(0)\sqrt{i^2+j^2}} = 0 . \quad (41)$$



$$R = 0.1, r = A_1, \sigma = 0.25$$

Figure 5: The trace of the topological evolution operator obtained from the BER approximation and directly from periodic orbits

This is derived directly from eqs. (28) and (29) but is indeed exact, thus demonstrating consistency of our approach. It allows determination of $h(0)$ within a fraction of a second of computer time. We find $h(0) = 1.9133307629\dots$. This value differs slightly from Berry's estimate [28] $h(0) \approx \sqrt{12/\pi} = 1.95441$.

Considering the one parameter family of weights $|\Lambda|^\tau$, we see that if $\tau > 0$ the leading zeros lie below the branch cut and the trace exhibit exponential behaviour. When $\tau = 0$ the leading zeros coincides with a branch point. When $\tau \leq 0$ the behaviour is entirely given by the branch cuts, i.e. it is governed by the tails in the free directions. This is usually referred to as a phase transition [29]. We have thus been able to give a nice analytical description of such a phase transition in terms of the zeta function; as the collision of the leading zero with a singularity.

5 Discussion

The aim has been to show that the BER approximation works, and works well for quite general weights w . We have not aimed at calculating more physically interesting properties, such as decay of correlations, fractal dimensions etc. This is of course a natural next step.

Regarding correlation functions one may, loosely, say that they behave very much like the trace of the evolution operator with unit weight. This is easily seen if the eigenvalues $\exp(ik_nt)$ are isolated. Then the correlation function $C(t)$ may (formally) be expanded as $C(t) = \sum_n c_n \exp(ik_nt)$. The coefficients c_n depend on the observables. The trace $\text{tr} \mathcal{L}^t = \sum_n \exp(ik_nt)$ may therefore be thought of as a archetype correlation function. The velocity autocorrelation function for the Sinai billiard (with continuous time) have been reported to decay as $1/t$ [30], which is the same decay as we found for the trace.

The BER approximation is in principle applicable to a wide range of bound ergodic Hamiltonian systems. The details have to be worked out for each separate case. For the Sinai billiard it turned out to be particularly simple. The reason for this is that we could make the approximation that the lengths l_s are constant over the phase space regions Ω_s . This was because the flow is

suspended in the sense that liminar segments of arbitrary lengths does not exist. For instance, there is no such length shorter than $1 - 2R$. This has to do with the presence of marginally stable orbits. A consequence of this is the periodicity of the cuts.

The situation is very different for e.g. the hyperbola billiard. In ref. [8] we found evidence that the zeta function ($\tau = 0$) for this system has a discrete spectrum of zeros and a single cut along the positive imaginary axis. We used cycle expansions to obtain this result, but since the zeta function is divergent we had to do some ad hoc manipulations. The expansion used exhibited some obvious similarities with the BER approximation worked out in this paper. The expansion consisted almost exclusively of cycles with only one laminar segment. To avoid (the most serious) divergence we selected out an infinite subsequence accumulating towards a pruned orbit and responsible for the branch cut. This sequence has as a counterpart the tails in the free directions of the Sinai billiard.

Perhaps the most powerful result in this article is the possibility of making asymptotic statements, cf eq (36), about periodic orbits. By varying the parameter τ one can obtain a wealth of such *periodic orbit sum rules*. This asymptotics is essential for doing semiclassical calculation on spectral fluctuations based on Gutzwiller's trace formula [31, 8]. We have therefore outlined how to do periodic orbit theory without a single periodic orbit.

Acknowledgements

I would like to thank Cecilia Kozma for a useful hint. This work was supported by the Swedish Natural Science Research Council (NFR) under contract no. F-FU 06420-303.

References

- [1] R. Artuso, E. Aurell and P. Cvitanović, *Nonlinearity* **3**, 325 and 361, (1990).
- [2] P. Cvitanović and B. Eckhardt, *Phys. Rev. Lett.* **63**, 823 (1989).
- [3] P. Cvitanović, P. E. Rosenqvist, G. Vattay, and H. H. Rugh, *A Fredholm Determinant for Semi-classical Quantization*
- [4] M. Pollicott, *Inv. Math.* **85**, 147, (1986).
- [5] D. Ruelle, *J. Stat.. Phys.* **44**, 281, (1986).
- [6] D. Ruelle, *J. Diff. Geom.* **25**, 99 and 117, (1986).
- [7] H. H. Rugh, *Nonlinearity* **5**, 1237, (1992).
- [8] P. Dahlqvist, *J. Phys. A* **27**, 763 (1994).
- [9] P. Dahlqvist, *J. Phys. A* **25**, 6265, (1992).
- [10] P. Cvitanović, *Physica D* **51**, 138 (1991).
- [11] K. Hansen, *Chaos* **2**, 71 (1991).
- [12] G. Tanner and D. Wintgen, *Classical and Semiclassical Zeta Functions in Terms of Transition Probabilities*, to appear in "Chaos, Solitons and Fractals".
- [13] P. Cvitanović and B. Eckhardt, *J. Phys. A* **24**, L237, (1991).
- [14] P. Cvitanović and G. Vattay, *Entire Fredholm Determinants for Evaluation of Semi-classical and Thermodynamical Spectra*
- [15] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, Springer, New York (1990).
- [16] A. Voros, *J. Phys. A* **21**, 685 (1988).
- [17] J. Robbins, *Phys. Rev. A* **40**, 2128 (1989).
- [18] B. Lauritzen, *Phys. Rev. A* **43**, 603 (1991).
- [19] P. Cvitanović and B. Eckhardt, *Nonlinearity* **6**, 277, (1993).
- [20] V. Baladi, J. P. Eckmann and D. Ruelle, *Nonlinearity* **2**, 119, (1989).
- [21] Y. G. Sinai, *Russ. Math. Surv.* **25**, 137, (1979).
- [22] J. Machta and B. Reinhold, *J. Stat.. Phys.* **42**, 949, (1986).
- [23] J-P. Bouchaud and P. L. Doussal, *J. Stat.. Phys.* **41**, 225, (1985).
- [24] Hamermesh, *Group Theory and its application to Physical Problems*, Reading MA: Addison-Wesley, (1962).
- [25] H. Rademacher, *Lectures on Elementary Number Theory*, Blaisdell Publishing Company, New York.
- [26] M. Sieber and F. Steiner, *Physica D* **44**, 248 (1990).
- [27] J. H. Hannay and A. M. Ozorio de Almeida, *J. Phys. A* **17**, 3429, (1984).
- [28] M. V. Berry, *Ann. Phys. (N.Y.)* **131**, 163, (1981).
- [29] R. Artuso, P. Cvitanović and B. G. Kenny, *Phys. Rev. A* **39**, 268, (1989).
- [30] B. Friedman and R. F. Martin, *Phys. Lett. A* **105**, 2 (1984).
- [31] M. V. Berry, *Proc. R. Soc. Lond. A* **400**, 229, (1985).